

# Trail County Community Health Profile

April 13, 2015



Stephen Pickard, MD  
North Dakota Department of Health

# TABLE OF CONTENTS

## POPULATION DATA

Chart 1: Population by Age Group.....	1
Chart 2: Pyramid Graph of Number of Residents by Census Year.....	1
Chart 3: Female Population and Percentage Female by Age.....	1
Chart 4: Decennial Population Change.....	1
Chart 5: Race.....	2
Chart 6: Household Populations.....	2
Chart 7: Marital Status of Persons Age 15 and Older.....	2
Chart 8: Educational Attainment Among Persons 25+.....	2
Chart 9: Persons with Disability.....	3
Chart 10: Income and Poverty Status by Age Group.....	3
Chart 11: Family Poverty and Childhood and Elderly Poverty.....	3
Chart 12: Age of Housing.....	3

## VITAL STATISTICS DATA

Birth and Deaths Definitions.....	4
Chart 13: Births.....	5
Chart 14: Child Deaths.....	5
Chart 15: Deaths and Age Adjusted Death Rate by Cause.....	5
Chart 16: Leading Causes of Death by Age Group for Traill County.....	6
Chart 17: Leading Causes of Death by Age Group for North Dakota.....	6

## ADULT BEHAVIORAL RISK FACTORS

Chart 18 –21:	
• Adult Behavioral Risk Factors Definition.....	7
• Alcohol.....	8
• Arthritis.....	8
• Asthma.....	8
• Body Weight.....	8
• Cancer.....	8
• Cardiovascular.....	9
• Cholesterol.....	9
• Chronic Lung Disease.....	9
• Colorectal Cancer.....	9
• Diabetes.....	9
• Fruits and Vegetables.....	9
• General Health.....	10
• Health Care Access.....	10
• Hypertension.....	10
• Immunization.....	10
• Injury.....	10
• Oral Health.....	11
• Physical Activity.....	11
• Tobacco.....	11
• Women’s Health.....	11

## CRIME

Chart 21: Traill County.....	12
Chart 22: North Dakota.....	12

## CHILD RISK

Chart 23: Child Indicators—Education.....	13
Chart 24: Child Indicators—Economic Health.....	13
Chart 25: Child Indicators—Families and Child Care.....	13
Chart 26: Child Indicators—Juvenile Justice.....	13


## POPULATION DATA

**1**

Population by Age Group, 2009-2013 ACS Estimates				
Traill County			North Dakota	
Age Group	Number	Percent	Number	Percent
0-9	963	11.9%	91,871	12.7%
10-19	1,075	13.2%	93,318	12.9%
20-29	973	12.0%	124,424	17.2%
30-39	852	10.5%	91,148	12.6%
40-49	1,001	12.3%	79,573	11.0%
50-59	1,197	14.7%	101,998	14.1%
60-69	909	11.2%	69,446	9.6%
70-79	576	7.1%	41,233	5.7%
80+	579	7.1%	31,829	4.4%
<b>Total</b>	<b>8,125</b>	<b>100.0%</b>	<b>724,840</b>	<b>100.0%</b>
0-17	1,786	22.0%	161,317	22.3%
65+	1,511	18.6%	102,722	14.2%

The Demographic Section of this report comes from the US Census Bureau ([www.census.gov](http://www.census.gov)). Most tables are derived either from the Census estimates for 2013 or from the Community Population Survey aggregated over a several year period. The table header describes the specific years from which the data is derived. The table showing percent population change uses census data from 2000 also. Tables present number of persons and percentages which in almost all circumstances represent the category specific percentage of all persons referenced by the table (e.g., percentage of persons age 15 and older who are married). Age specific poverty rates represent the percentage of each age group which is in poverty (e.g., percentage of children under five years in poverty).

**2**



**3**

Female Population and Percentage Female by Age, 2009-2013 ACS Estimates				
Traill County			North Dakota	
Age Group	Number	Percent	Number	Percent
0-9	453	47.0%	45,342	49.4%
10-19	534	49.7%	44,988	48.2%
20-29	449	46.1%	55,615	44.7%
30-39	425	49.9%	41,092	45.1%
40-49	478	47.8%	39,675	49.9%
50-59	557	46.5%	50,656	49.7%
60-69	477	52.5%	34,361	49.5%
70-79	301	52.3%	21,254	51.5%
80+	381	65.8%	20,546	64.5%
<b>Total</b>	<b>4,055</b>	<b>49.9%</b>	<b>353,530</b>	<b>48.8%</b>
0-17	865	48.4%	78,484	48.7%
65+	882	58.4%	50,201	48.9%

**4**

Decennial Population Change, 1990 to 2000, 2000 to 2010				
Census	Traill County	10 Year Change	North Dakota	10 Year Change
1990	8,752	(%)	638,800	(%)
2000	8,477	-3.1%	642,200	0.5%
2010	8,121	-4.2%	672,591	4.7%

## POPULATION DATA

**5**

Race, 2009-2013 ACS Estimates				
Race	Traill County		North Dakota	
	Number	Percentage	Number	Percentage
Total	8,125	100.0%	723,393	100.0%
White	7,831	96.4%	643,478	89.0%
Black	43	0.5%	10,827	1.5%
Am. Indian	82	1.0%	40,214	5.6%
Asian	11	0.1%	9,096	1.3%
Pac. Islander	0	0.0%	371	0.1%
Other	61	0.8%	4,620	0.6%
Multirace	97	1.2%	29,574	4.1%

**6**

Household Populations, 2011-2013 ACS Three Year Estimates				
	Traill County		North Dakota	
	Number	Percent	Number	Percent
Total	8,125	100.0%	703,203	100.0%
In Family Households	6,194	76.2%	530,615	75.5%
In Non-Family Households	1,591	19.6%	146,330	20.8%
Total In Households	7,785	95.8%	676,945	96.3%
Institutionalized*	168	2.1%	9,675	1.4%
Non-institutionalized*	172	2.1%	16,583	2.4%
Total in Group Quarters	340	4.2%	26,258	3.7%

**7**

Marital Status of Persons Age 15 and Older, 2009-2013 ACS Estimates				
Marital Status	Traill County		North Dakota	
	Number	Percent	Number	Percent
Total Age 15+	6,677	100.0%	561,346	100.0%
Never Married	1,823	27.3%	177,385	31.6%
Now Married	3,559	53.3%	291,900	52.0%
Separated	47	0.7%	5,052	0.9%
Widowed	668	10.0%	33,681	6.0%
Divorced	581	8.7%	53,328	9.5%

**8**

Educational Attainment Among Persons 25+, 2009-2013 ACS Estimates				
Education	Traill County		North Dakota	
	Number	Percent	Number	Percent
Total	5,519	100.0%	457,771	100.0%
Less than 9th Grade	243	4.4%	19,226	4.2%
Some High School	281	5.1%	21,057	4.6%
High school or GRE	1,391	25.2%	125,429	27.4%
Some College / Asso. Degree	2,009	36.4%	168,002	36.7%
Bachelor's degree	1,269	23.0%	89,723	19.6%
Post Graduate Degree	326	5.9%	34,791	7.6%

## POPULATION DATA

9

Persons with Disability, 2009-2013 ACS Estimates					
Group	Traill County		North Dakota		
	Number	Percent	Number	Percent	
Total	7,984	100.0%	688,158	100.0%	
Any Disability	855	10.7%	72,762	10.6%	
No Disability	7,129	89.3%	615,396	89.4%	
Self Care Disability	156	2.0%	11,274	1.6%	
0-17 with any disability	66	3.7%	4,677	2.9%	
18-64 with any disability	402	8.3%	35,931	7.8%	
65+ with any disability	387	28.1%	32,154	31.3%	

10

Income and Poverty Status by Age Group, 2009-2013 ACS Estimates				
	Traill County		North Dakota	
	Number	Percent	Number	Percent
Median Household Income	\$48,767		\$54,920	
Per Capita Income	\$27,096		\$30,436	
	Number	Percent	Number	Percent
Below Poverty Level	828	10.7%	80,644	11.9%
Under 5 years	21	17.4%	7,714	16.9%
5 to 11 years	125	17.4%	7,944	13.2%
12 to 17 years	48	8.3%	5,776	11.8%
18 to 64 years	427	9.2%	49,568	11.6%
65 to 74 years	52	7.6%	3,448	7.0%
75 years and over	110	15.8%	6,194	13.7%
Total Known Children in Poverty	194	10.9%	21,434	13.3%
Total Known Age 65+ in Poverty	162	10.7%	9,642	9.4%

11

Family Poverty, 2009-2013 ACS Estimates				
	Traill County		North Dakota	
	Number	Percent*	Number	Percent*
Total Families	1,964	100.0%	176,378	100.0%
Families in Poverty	132	6.7%	13,052	7.4%
Families with Own Children	842	42.9%	80,964	45.9%
Families with Own Children in Poverty	103	5.2%	10,121	5.7%
Families with Own Children and Female Parent Only	144	7.3%	16,716	9.5%
Families with Own Children and Female Parent Only in Poverty	80	4.1%	6,452	3.7%

12

Age of Housing, 2009-2013 ACS Estimates				
	Traill County		North Dakota	
	Number	Percent	Number	Percent
Housing units: Total	3,778	100.0%	329,970	100.0%
1980 and Later	815	21.6%	128,111	38.8%
1970 to 1979	748	19.8%	66,396	20.1%
Prior to 1970	2,215	58.6%	135,463	41.1%

## Vital Statistics Data

### BIRTHS AND DEATHS DEFINITIONS

Vital Statistics Data comes from the birth and death records collected by the State of North Dakota aggregated over a five year period. All births and deaths represent the county of residence not the county of occurrence. The number of events is blocked if fewer than five. Formulas for calculating rates and ratios are as follows:

**Birth Rate** = Resident live births divided by the total resident population x 1000.

**Pregnancies** = Live births + Fetal deaths + Induced termination of pregnancy.

**Pregnancy Rate** = Total pregnancies divided by the total resident population x 1000.

**Fertility Rate** = Resident live births divided by female population (age 15-44) x 1000.

**Teenage Birth Rate** = Teenage births (age <20) divided by female teen population x 1000.

**Teenage Pregnancy Rate** =  
Teenage pregnancies (age <20) divided by female teen population x 1000.

**Out of Wedlock Live Birth Ratio** =  
Resident OOW live births divided by total resident live births x 1000.

**Out of Wedlock Pregnancy Ratio** =  
Resident OOW pregnancies divided by total pregnancies x 1000.

**Low Weight Ratio** =  
Low weight births (birth weight < 2500 grams) divided by total resident live births x 1000.

**Infant Death Ratio** = Number of infant deaths divided by the total resident live births x 1000.

**Childhood & Adolescent Deaths** = Deaths to individuals 1 - 19 years of age.

**Childhood and Adolescent Death Rate** =  
Number of resident deaths (age 1 - 19) divided by population (age 1 - 19) x 100,000.

**Crude Death Rate** = Death events divided by population x 100,000.

**Age-Adjusted Death Rate** = Death events with age specific adjustments x 100,000 population.

## Vital Statistics Data

### BIRTHS AND DEATHS

13

Births, 2009-2013	Traill County		North Dakota	
	Number	Rate or Ratio	Number	Rate or Ratio
Live Births and Rate	474	12	47,959	14
Pregnancies and Rate	502	12	52,505	15
Fertility Rate		70		72
Teen Births and Rate	23	11	2,118	12
Teen Pregnancies and Rate	29	14	3,725	21
Out of Wedlock Births and Ratio	131	276	15,686	327
Out of Wedlock Preg and Ratio	156	311	19,436	370
Low Birth Weight Birth and Ratio	8	17	3,078	64

14

Child Deaths, 2009-2013	Traill County		North Dakota	
	Number	Rate or Ratio	Number	Rate or Ratio
Infant Deaths and Ratio	*	NA	286	6
Child and Adolescent Deaths and Rate	*	NA	270	32
Total Deaths and Crude Rate	448	1,103	29,616	866

15

Deaths and Age Adjusted Death Rate by Cause, 2009-2013	Traill County		North Dakota	
	Number (Adj. Rate)	Number (Adj. Rate)	Number (Adj. Rate)	Number (Adj. Rate)
All Causes	448 (661)		29,581 (702)	
Heart Disease	133 (169)		6,762 (154)	
Cancer	89 (154)		6,315 (156)	
Stroke	36 (44)		1,664 (37)	
Alzheimers Disease	31 (37)		2,189 (45)	
COPD	24 (36)		1,707 (41)	
Unintentional Injury	17 (34)		1,625 (44)	
Diabetes Mellitus	9 (13)		1,022 (24)	
Pneumonia and Influenza	9 (13)		682 (15)	
Cirrhosis	*		394 (11)	
Suicide	*		551 (16)	

## Vital Statistics Data

### BIRTHS AND DEATHS

16


Leading Causes of Death by Age Group for Traill County, 2009-2013			
Age	1	2	3
0-4	Congenital Anomaly		
5-14			
15-24	Unintentional Injury		
25-34	Unintentional Injury		
35-44	Cancer	Suicide	Unintentional Injury Cirrhosis
45-54	Cancer 6	Heart	Unintentional Injury
55-64	Cancer 10	Heart 7	COPD
65-74	Cancer 19	Heart 12	Stroke 5
75-84	Cancer 32	Heart 31	COPD 11
85+	Heart 81	Stroke 24	Alzheimer's Dz 21

17

Leading Causes of Death by Age Group for North Dakota, 2006-2010			
Age	1	2	3
0-4	Congenital Anomaly 69	Prematurity 44	SIDS 40
5-14	Unintentional Injury 26	Cancer 10	Congenital Anomaly 6
15-24	Unintentional Injury 184	Suicide 109	Cancer 20
25-34	Unintentional Injury 166	Suicide 91	Heart 32
35-44	Unintentional Injury 173	Heart 94	Cancer 88
45-54	Cancer 493	Heart 335	Unintentional Injury 194
55-64	Cancer 1001	Heart 579	Unintentional Injury 137
65-74	Cancer 1562	Heart 843	COPD 313
75-84	Cancer 1992	Heart 1797	COPD 626
85+	Heart 3421	Alzheimer's Dz 1391	Cancer 1352

## ADULT BEHAVIORAL RISK FACTORS DEFINITION

The following three pages represent data received from the Adult Behavioral Risk Factor Surveillance Survey. The Adult Behavioral Risk Factor data are derived from aggregated data (the number of years specified is in the table) continuously collected by telephone survey from persons 18 years and older. All data is self-reported data. Numbers given are point estimate percentages followed by 95% confidence intervals. Statistical significance can be determined by comparing confidence intervals between two geographic areas. To be statistically significant, confidence intervals may not overlap. For example the confidence intervals 9.3 (8.3-10.2) and 10.8 (10.0-11.6) overlap (see picture below) so the difference between the two numbers is not statistically significant. That means that substantial uncertainty remains whether the apparent difference is due to chance alone (due to sampling variation) rather than representing a true difference in the prevalence of the condition in the two populations. The less they overlap, the more likely it is that the point estimates represent truly different prevalences in the two populations.



## ADULT BEHAVIORAL RISK FACTORS, 2011-2013

18

	<b>ALCOHOL</b>	<b>Trall and Steele 2011-2013</b>	<b>Trall, Steele, Sargent, Ransom 2011-2013</b>	<b>North Dakota 2011-2013</b>
Binge Drinking	Respondents who reported binge drinking (5 drinks for men, 4 drinks for women) one or more times in the past 30 days.	20.3 (12.4-28.3)	24.3 (18.8-29.9)	24.2 (23.2-25.1)
Heavy Drinking	Respondents who reported heavy drinking (more than 2 drinks per day for men, more than 1 drink per day for women) during the past 30 days	NA	7.5 ( 4.3-10.7)	6.8 (6.2-7.4)
Drunk Driving	Respondents who reported driving when they had too much to drink one or more times during the past 30 days	NA	NA	1.9 (1.5-2.4)
<b>ARTHRITIS</b>				
Doctor Diagnosed Arthritis	Respondents who reported ever have been told by a doctor or other health professional that they had some form of arthritis.	24.2 (18.5-29.9)	28.8 (24.4-33.3)	25.1 (24.3-25.9)
Activity Limitation Due to Arthritis	Respondents who reported being limited in any usual activities because of arthritis or joint symptoms.	NA	9.5 ( 6.6-12.4)	8.3 (7.8-8.8)
<b>ASTHMA</b>				
Ever Asthma	Respondents who reported ever having been told by a doctor, nurse or other health professional that they had asthma.	10.0 ( 5.3-14.7)	10.6 ( 7.1-14.2)	11.5 (10.7-12.2)
Current Asthma	Respondents who reported ever having been told by a doctor, nurse or other health professional that they had asthma and who still have asthma.	NA	8.2 ( 4.9-11.6)	8.1 (7.5-8.7)
<b>BODY WEIGHT</b>				
Overweight But Not Obese	Respondents with a body mass index greater than or equal to 25 but less than 30 (overweight)	39.0 (30.2-47.9)	37.8 (32.0-43.7)	36.7 (35.7-37.7)
Obese	Respondents with a body mass index greater than or equal to 30 (obese)	37.2 (28.1-46.2)	33.8 (28.1-39.5)	29.4 (28.5-30.4)
Overweight or Obese	Respondents with a body mass index greater than or equal to 25 (overweight or obese)	76.2 (68.2-84.2)	71.6 (65.9-77.3)	66.1 (65.1-67.2)
<b>CANCER</b>				
Ever Cancer	Respondents who reported ever having been told by a doctor, nurse or other health professional that they had cancer (excluding skin cancer).	9.0 ( 5.3-12.7)	7.5 ( 5.2- 9.9)	6.4 (6.0-6.8)

## ADULT BEHAVIORAL RISK FACTORS, 2011-2013

**19**

	<b>CARDIOVASCULAR</b>	<b>Trall and Steele 2011-2013</b>	<b>Trall, Steele, Sargent, Ransom 2011-2013</b>	<b>North Dakota 2011-2013</b>
Heart Attack	Respondents who reported ever having been told by a doctor, nurse or other health care professional that they had a heart attack.	NA	4.0 ( 2.5- 5.5)	4.3 (3.9-4.6)
Angina	Respondents who reported ever having been told by a doctor, nurse or other health care professional that they had angina.	NA	3.3 ( 1.9- 4.7)	4.1 (3.7-4.4)
Stroke	Respondents who reported ever having been told by a doctor, nurse or other health care professional that they had a stroke.	NA	NA	2.3 (2.0-2.5)
Cardiovascular Disease	Respondents who reported ever having been told by a doctor, nurse or other health care professional that they had any of the following: heart attack, angina or stroke.	6.2 ( 3.3- 9.1)	6.5 ( 4.5- 8.4)	7.6 (7.1-8.0)
<b>CHOLESTEROL</b>				
Never Cholesterol Test	Respondents who reported never having a cholesterol test	NA	19.0 (12.6-25.4)	22.3 (21.1-23.4)
No Cholesterol Test in Past 5 Years	Respondents who reported never having a cholesterol test in the past five years	NA	24.6 (18.0-31.1)	26.7 (25.5-27.9)
High Cholesterol	Respondents who reported that they had ever been told by a doctor, nurse or other health professional that they had high cholesterol.	40.3 (30.2-50.4)	40.3 (33.8-46.9)	36.6 (35.4-37.8)
<b>CHRONIC LUNG DISEASE</b>				
COPD	Respondents who have ever been told by a doctor, nurse or other health professional ever told you that they have COPD (chronic obstructive pulmonary disease), emphysema, or chronic bronchitis?	NA	5.7 ( 3.6- 7.9)	4.6 (4.2-5.0)
<b>COLORECTAL CANCER</b>				
Fecal Occult Blood	Respondents age 50 and older who reported not having a fecal occult blood test in the past two years.	90.0 (81.6-98.4)	81.5 (73.2-89.9)	86.2 (84.8-87.6)
Never Sigmoidoscopy	Respondents age 50 and older who reported never having had a sigmoidoscopy or colonoscopy	NA	45.2 (33.4-56.9)	38.0 (35.9-40.2)
Up to date for Colorectal Screening	Respondents age 50 and older who are up to date according to recommended screening guidelines for colorectal screening	NA	53.8 (42.1-65.4)	59.1 (56.9-61.3)
<b>DIABETES</b>				
Diabetes Diagnosis	Respondents who reported ever having been told by a doctor that they had diabetes.	10.5 ( 6.1-14.8)	9.9 ( 7.0-12.8)	8.5 (8.0-9.0)
<b>FRUITS AND VEGETABLES</b>				
Five Fruits and Vegetables	Respondents who reported that they do not usually eat 5 fruits and vegetables per day	84.4 (76.2-92.7)	87.5 (82.8-92.2)	85.9 (85.0-86.7)

## ADULT BEHAVIORAL RISK FACTORS, 2011-2013

20

	GENERAL HEALTH	Traill and Steele 2011-2013	Traill, Steele, Sargent, Ransom 2011-2013	North Dakota 2011-2013
Fair or Poor Health	Respondents who reported that their general health was fair or poor	14.0 ( 8.8-19.2)	11.8 ( 8.6-14.9)	14.0 (13.3-14.7)
Poor physical Health	Respondents who reported they had 8 or more days in the last 30 when their physical health was not good	10.5 ( 5.6-15.4)	9.1 ( 5.8-12.4)	11.6 (11.0-12.3)
Poor Mental Health	Respondents who reported they had 8 or more days in the last 30 when their mental health was not good	11.2 ( 4.9-17.4)	10.3 ( 6.2-14.4)	10.8 (10.2-11.5)
Activity Limitation Due to Poor Health	Respondents who reported they had 8 or more days in the last 30 when poor physical or mental health kept them from doing their usual activities.	NA	NA	7.1 (6.6-7.6)
Any Activity Limitation	Respondents who reported being limited in any way due to physical, mental or emotional problem.	17.3 (11.7-22.9)	15.2 (11.7-18.7)	17.9 (17.2-18.7)
<b>HEALTH CARE ACCESS</b>				
Health Insurance	Respondents who reported not having any form or health care coverage	NA	9.0 ( 5.5-12.4)	12.4 (11.6-13.1)
Access Limited by Cost	Respondents who reported needing to see a doctor during the past 12 months but could not due to cost.	NA	NA	8.1 (7.4-8.7)
No Personal Provider	Respondents who reported that they did not have one person they consider to be their personal doctor or health care provider.	20.7 (12.8-28.6)	20.4 (15.5-25.4)	26.0 (25.0-26.9)
<b>HYPERTENSION</b>				
High Blood Pressure	Respondents who reported ever having been told by a doctor, nurse or other health professional that they had high blood pressure.	24.0 (16.7-31.2)	28.9 (23.5-34.3)	29.5 (28.6-30.6)
<b>IMMUNIZATION</b>				
Influenza Vaccine	Respondents age 65 and older who reported that they did not have a flu shot in the past year	24.2 (14.5-33.9)	32.5 (24.2-40.9)	40.3 (38.6-42.0)
Pneumococcal Vaccine	Respondents age 65 or older who reported never having had a pneumonia shot.	NA	28.2 (19.6-36.8)	29.4 (27.8-31.0)
<b>INJURY</b>				
Fall	Respondents 45 years and older who reported that they had fallen in the past 12 months	NA	28.2 (19.0-37.5)	27.8 (25.9-29.6)
Seat Belt	Respondents who reported not always wearing their seatbelt	45.7 (37.1-54.3)	47.8 (41.9-53.7)	32.6 (31.6-33.6)

## ADULT BEHAVIORAL RISK FACTORS, 2011-2013

21

	ORAL HEALTH	Traill and Steele 2011-2013	Traill, Steele, Sargent, Ransom 2011-2013	North Dakota 2011-2013
Dental Visit	Respondents who reported that they have not had a dental visit in the past year	NA	NA	32.8 (31.0-34.7)
Tooth Loss	Respondents who reported they ever had a permanent tooth extracted.	NA	NA	43.2 (41.4-45.0)
<b>PHYSICAL ACTIVITY</b>				
Recommend Physical Activity	Respondents who reported that they did not get the recommended amount of physical activity	NA	51.7 (44.6-58.8)	53.8 (52.6-55.1)
Inactive	Respondents reporting little or no physical activity	24.8 (15.8-33.9)	29.7 (23.4-36.0)	30.4 (29.3-31.6)
<b>TOBACCO</b>				
Current Smoking	Respondents who reported that they smoked every day or some days	17.4 (10.4-24.5)	18.0 (13.4-22.5)	21.6 (20.6-22.5)
<b>WOMEN'S HEALTH</b>				
Pap Smear	Women 18 and older who reported that they have not had a pap smear in the past three years	NA	NA	23.7 (20.9-26.5)
Mammogram Age 40+	Women 40 and older who reported that they have not had a mammogram in the past two years	NA	NA	26.4 (24.0-28.9)

## CRIME

**21**

<b>Trail County</b>							
	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>	<b>2013</b>	<b>5 year</b>	<b>5-Year Rate</b>
Murder	0	0	0	0	0	0	0.0
Rape	0	0	1	0	1	2	4.8
Robbery	1	0	0	0	0	1	2.4
Aggrav. Assault	0	0	2	2	2	6	14.5
<b>Violent crime</b>	<b>1</b>	<b>0</b>	<b>3</b>	<b>2</b>	<b>3</b>	<b>9</b>	<b>21.8</b>
Burglary	9	16	8	16	11	60	145.3
Larceny	26	27	36	42	26	157	380.2
Motor vehicle theft	6	8	2	3	9	28	67.8
<b>Property crime</b>	<b>41</b>	<b>51</b>	<b>46</b>	<b>61</b>	<b>46</b>	<b>245</b>	<b>593.4</b>
<b>Total</b>	<b>42</b>	<b>51</b>	<b>49</b>	<b>63</b>	<b>49</b>	<b>254</b>	<b>615.2</b>

**22**

<b>North Dakota</b>							
	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>	<b>2013</b>	<b>5 year</b>	<b>5-Year Rate</b>
Murder	15	11	15	20	14	75	2.2
Rape	206	222	207	243	237	1,115	32.6
Robbery	102	85	91	117	151	546	16.0
Aggrav. Assault	795	847	1,040	1,071	1,156	4,909	143.6
<b>Violent crime</b>	<b>1,118</b>	<b>1,165</b>	<b>1,353</b>	<b>1,451</b>	<b>1,558</b>	<b>6,645</b>	<b>194.3</b>
Burglary	2,180	1,826	2,227	2,200	2,656	11,089	324.3
Larceny	8,699	8,673	9,344	10,184	10,243	47,143	1378.6
Motor vehicle theft	854	825	763	854	1,228	4,524	132.3
<b>Property crime</b>	<b>11,733</b>	<b>11,324</b>	<b>12,334</b>	<b>13,238</b>	<b>14,127</b>	<b>62,756</b>	<b>1835.2</b>
<b>Total</b>	<b>12,851</b>	<b>12,489</b>	<b>13,687</b>	<b>14,689</b>	<b>15,685</b>	<b>69,401</b>	<b>2029.5</b>

## CHILD HEALTH INDICATORS

**23**

Child Indicators: Education 2013	Traill County		North Dakota	
	Number	Percent	Number	Percent
Enrolled in Special Education Ages 3-21 (Number and percent of total school enrollment)	152	11.0%	13,399	13.0%
High School Dropouts (Dropouts per 100 persons Grades 9-12)*	5	1.2%	888	2.8%
Average Expenditure per Student in Public School	\$9,527		\$10,964	

**24**

Child Indicators: Economic Health 2013	Traill County		North Dakota	
	Number	Percent	Number	Percent
TANF Recipients Ages 0-19 (Percentage of persons ages 0-19)	27	1.3%	5,358	2.9%
Food Stamp (SNAP) Recipients Ages 0-18 (Percentage of all children ages 0-19)	378	19%	37,826	23%
Children Receiving Free or Reduced Price Lunch (Percentage of total school enrollment)	341	25%	34,381	31%
Medicaid Recipients Ages 0-20 (Percentage of all persons ages 0-20)*	528	24%	53,814	27%
Children Ages 0-17 Living in Extreme Poverty (Percentage of children 0-17 for whom poverty is determined)	131	7.4%	10,114	6.2%
Median Income for Families with Children Ages 0-17 (Percentage of all women with children ages 0-17)	\$70,208		\$70,530	

**25**

Child Indicators: Families and Child Care 2013	Traill County		North Dakota	
	Number	Percent	Number	Percent
Mothers in Labor Force with a Child Ages 0-17 (Percentage of all mothers with a child ages 0-17)	726	85%	57,908	81%
Children Ages 0-17 Living in a Single Parent Family (Percentage of all children ages 0-17)	323	18%	34,591	23%
Children in Foster Care (Percentage of children ages 0-18)*	23	1.2%	2,019	1.2%
Children Ages 0-17 with Suspected Child Abuse or Neglect (Percentage of children 0-17)	36	2.0%	6,170	4.0%
Children Ages 0-17 Impact by Domestic Violence (Percentage of all children ages 0-17)**	14	0.8%	4,739	2.9%
Births to Mothers with Inadequate Prenatal Care (Percentage of Total Births)	8	8.0%	508	4.8%

\*\*2010

**26**

Child Indicators: Juvenile Justice 2013	Traill County		North Dakota	
	Number	Percent	Number	Percent
Children Ages 10-17 Referred to Juvenile Court (Percentage of all children ages 0-17)	13	1.6%	3,789	5.8%
Offense Against Person Juvenile Court Referral (Percentage of total juvenile court referral)	1	6.0%	689	9.8%
Alcohol-Related Juvenile Court Referral (Percentage of all juvenile court referrals)*	5	30%	909	13%