


North Dakota Mosquito Surveillance 2009 Program



NORTH DAKOTA
DEPARTMENT *of* HEALTH



Division
of
Laboratory Services

Collaborators

Michelle Henke
Mosquito Surveillance Program Coordinator

Sarah Iverson
Summer Mosquito Analyst

Abby Coleman
Summer Mosquito Analyst

Myra Kosse
Division of Laboratory Services Director

Mike Trythall
Arbovirus Program Supervisor



North Dakota Department of Health
Division of Laboratory Services
2635 East Main Avenue
Bismarck, North Dakota 58506-5520

2009 North Dakota Mosquito Surveillance Program's Mission

Through mosquito collection and speciation, the North Dakota Department of Health (NDDoH) monitors the risk of infection from arboviral encephalitides that are known to occur in this region. The North Dakota Mosquito Surveillance team focuses activities on *Culex tarsalis*, monitoring for increased numbers in the New Jersey mosquito trap network and viral identification using the CDC miniature light mosquito trap network. Should mosquito populations reach significant levels or arbovirus activity is detected, appropriate recommendations for mosquito population control will be issued by the NDDoH to the vector control districts.

Table of Contents

North Dakota Mosquito Surveillance Program Background	4
New Jersey Mosquito Trap Network	
Operator List	5
2009 Network State Map	7
Network Information	8
2009 Species Comparison Graph	8
2009 Network Mosquito Count Results	9
Arbovirus Information	10
Mosquito Surveillance Risk Assessment Chart	12
Appendix A: 2009 New Jersey Mosquito Trap Data Analysis	
Introduction	13
Graphs	15
Appendix B: 1994-2009 Weekly New Jersey Mosquito Trap Counts Comparison	
Introduction	25
Trap Count Graphs	25
References	33

North Dakota Mosquito Surveillance Program Background

Since 1975, the North Dakota Department of Health has monitored the mosquito populations throughout the state. The Mosquito Surveillance Program traditionally has been activated following arboviral outbreaks or flooding incidences in various locations statewide.

The program was first initiated in 1975 following an outbreak of western equine encephalitis (WEE) and St. Louis encephalitis (SLE) in the United States. In 1977, the program was officially formed under the title *North Dakota Arboviral Encephalitis Surveillance Program* and housed with the Division of Environmental Sanitation and Food Protection. This program was responsible for equine and human arbovirus surveillance until 1989.

The program was reinstated under the name *North Dakota Mosquito Surveillance Program* in 1994 in response to flooding of the Red River in 1993. This program was operated by the Division of Microbiology until 1997.

In 2000, the *North Dakota Mosquito Surveillance Program* was reinstated in response to the 1999 West Nile virus (WNV) outbreak in New York. In 2002, North Dakota had its first confirmed human cases of WNV, as well as detectable virus through laboratory testing in birds, horses and mosquitoes.

The 2003 program was expanded from 50 New Jersey mosquito traps to a network of 87 traps and 18 CDC miniature light mosquito traps. These enhancements provided network coverage statewide. The 2004 program further expanded the trap network to include 94 New Jersey mosquito traps and 33 CDC miniature light mosquito traps. A video also was produced to aid in trap-placement training. The 2005 program was further expanded to 103 New Jersey mosquito traps and 39 CDC miniature light mosquito traps. The program for 2006 had 100 New Jersey traps in operation, with at least one in each county. The dry conditions during the 2006 season kept the mosquito numbers low when compared to other seasons, and it was decided to postpone any live trapping. In 2007, we had 97 New Jersey traps in use. In July, live trapping was initiated at nine locations in Grand Forks, at two locations on the grounds of the laboratory, and at one location set up by the city of Bismarck due to increased *Culex tarsalis* numbers in the state. Four out of 17 pools collected at the laboratory tested positive for West Nile virus. All 14 pools collected by the city of Grand Forks and the five pools collected by the city of Bismarck were negative. In 2008, trappers across the state maintained a New Jersey light trap network of 92 traps. Live trapping was not implemented in 2008 by the Division of Laboratory Services - Microbiology. For 2009, there were 91 total New Jersey light traps in operation. Live trapping was not implemented.

Information about West Nile virus in North Dakota is available at www.ndhealth.gov/wnv.

New Jersey Mosquito Trap Network

The New Jersey mosquito trap network monitors mosquito populations throughout the state. By identifying mosquito populations known to be competent encephalitis vectors, the information from the network is used to determine the threat of mosquito-borne encephalitis in various regions of the state.

Thank you to the following New Jersey mosquito trap operators whose dedication and commitment to the North Dakota Department of Health Mosquito Surveillance Program made the 2009 program a success!


*Indicates State Park ** Indicates National Wildlife Refuge

Location	Trapper	Location	Trapper	Location	Trapper
Almont	Idar Handegard	Crosby	Dennis Lampert	Grand Forks	Todd Hanson
Amidon	Lane Smith	Cross Ranch*	Eric Lang	Hannover	Anton Sattler
Arrowwood**	Paulette Scherr	Deep Water Creek Bay	Kerry Hartman	Harvey	Loren Stolz
Ashley	Marcus Lynn	Devils Lake	Myron Asleson Leroy Axdahl	Hazelton	Bev Voller
Beach	Bill & Jeanie Larson	Dickinson	Skip Rapp Denny Smith	Hazen	Anton Sattler
Beaver Lake*	James Loken	Edmore	Roger Nygard	Hebron	Chad Stern Lance Elmer
Belfield	Susan Heck	Enderlin	Rick Gillund	Hettinger	Julie Kramlich
Beulah	Anton Sattler	Fargo	Brady Schribner	Hillsboro	Jim Anderson
Bismarck	Mel Fischer	Finley	Diane Jacobson	Icelandic*	Justin Robinson
Bottineau	Keith Fulsebakke	Forman	Colleen Sundquist	Indian Hills Recreational Area*	Byron and Tolly Holtan
Bowbells	Petter Willyard	Fort Ransom*	John Kwapinski	J.Clark Saylor **	Gary Erickson
Bowman	Andrea Bowman	Fort Stevenson*	Richard Messerly; Ryan Nelson	Jamestown	Steve Reidburn
Camp Grassick, Dawson	Dan Mimnaugh	Fort Totten	Hilda Garcia	Lake Metigoshe*	Larry Hagen
Cando	Duane Holien	Fort Yates	Bill Sherwood	Lake Sakakawea*	Keith Orth
Carrington	Shaunette Koenig; Jackie Schroeder	Ft. Abraham Lincoln*	Dan Schelske	Lamoure	Tony Hanson
Casselton	Camille Paaverud	Grafton	Mike Huska	Langdon	Rob Gilseth
Cooperstown	Farrah Saxberg	Grahams Island*	Dick Horner Trevor Retterath	Lewis & Clark*	Alan Larson

New Jersey Mosquito Trap Network Continued

Location	Trapper		Location	Trapper		Location	Trapper
Lisbon	Randy Seelig		Mott	Kim Kibbel		Tioga	Don Zacharias Kirk Odegard
Maddock	Frank Mosser		Napoleon	Sheldon Gerhardt		Towner	Nancy Bryn
Mandan	Dick Bechtel		New Rockford	George Ritzke		Turtle River*	Steve Best
Manning	Kevin Pavlish		Oakes	Robert Schaefer		Upper Souris**	Thomas Pabian
Martin	Bonita Kluck		Pembina	Kathy Johnson		Valley City	Jeff Differding
McVille	Julie Ferry		Rolla	Scott Hanson		Wahpeton	Randy Nelson
Medora	Bill Whitworth		Rugby	Deb Schiff		Washburn	Sandy Birst
Minot	Jim Heckman		Ryder	Jody Reinsch		Watford City	Kristen Gravos
Mohall	Tammy Aberle		Stanley	Jim Hennessy		Williston	Mike Melius Dan Saint

2009 New Jersey Mosquito Trap Surveillance Sites & Regions




New Jersey Mosquito Trap Network Information

In 2009, the New Jersey mosquito trap network had a total of 91 traps across North Dakota. There were 12 in state parks and three in national wildlife refuges. Two New Jersey mosquito traps were located in each urban area with a population greater than 5,000 citizens.

At the beginning of mosquito season, usually Memorial Day, the New Jersey mosquito trap operator installs a trap in a suitable location. Using a programmable timer, the trap is set to operate from dusk to dawn seven nights a week. At the end of the seven-day period, the trap contents are collected and sent to the North Dakota Department of Health, Division of Laboratory Services - Microbiology in Bismarck for counting and speciation. This process is repeated weekly until Labor Day.

At the Division of Laboratory Services - Microbiology, mosquito surveillance personnel sort the mosquitoes by sex and genera. Since male mosquitoes do not bite, they are of little health concern. However, their numbers are monitored because male mosquitoes hatch first, and increased numbers may indicate a future female mosquito population boom. The female mosquitoes are separated into four genera: *Anopheles*, *Aedes*, *Culex* and *Culiseta*. These genera are then enumerated.

- *Anopheles* is associated with malaria and West Nile virus.
- *Aedes* is associated with illnesses such as canine heartworm, LaCrosse encephalitis (LCE), eastern equine encephalitis (EEE), western equine encephalitis (WEE), California encephalitis (CAE), and West Nile virus (WNV). Although *Aedes vexans* has been shown to be capable of laboratory transmission of WNV, its mammalian feeding preferences decrease its potential as an enzootic vector for WNV.
- *Culex* is the mosquito of greatest public health concern in North Dakota, since all species are competent vectors of SLE, WEE and WNV. The species most commonly associated with encephalitis in North Dakota is *Culex tarsalis*, a principal arbovirus vector in rural agricultural ecosystems.
- *Culisetas* are monitored due to its association with eastern equine encephalitis.



2009 New Jersey Mosquito Trap Count Totals by Week – Counties

Week of	Male	Female									Total mosquitoes	Sites reporting out of 76
		Anopheles	Aedes	Aedes vexans	Culex	Culex tarsalis	Culex salinarius	Culiseta	Other	Total female		
1	628	26	1,512	0	131	5	0	46	0	1,720	2,348	47
June 1-8	645	54	1,446	0	32	1	0	58	0	1,591	2,236	62
June 9-14	3,440	249	2,606	774	278	1	4	171	3	4,086	7,526	67
June 15-21	5,459	216	8,597	1,270	978	188	12	703	0	11,964	17,423	67
June 22-28	11,407	222	9,242	884	398	488	10	1,586	0	12,830	24,237	64
5	12,203	129	21,153	16	119	1,114	0	2,038	0	24,569	36,772	67
July 6-12	10,217	159	14,377	1,041	266	1,976	1	2,389	0	20,209	30,426	65
July 13-19	30,186	17	34,261	561	26	2,895	1	3,723	0	41,484	71,670	65
July 20-26	22,882	226	29,669	4,213	729	3,327	0	2,113	0	40,277	63,159	68
July 27- August	11,656	170	7,450	1,047	156	1,457	17	859	0	11,156	22,812	65
August 3-9	2,809	124	4,518	435	97	573	17	461	0	6,225	9,034	61
August 10-16	7,104	174	9,392	476	228	1,629	8	848	0	12,755	19,859	65
August 17-24	595	17	1,060	15	837	125	0	18	0	2,072	2,667	57
August 24-31	883	0	1,779	88	536	45	0	21	0	2,469	3,352	47
August 31- September 7	1,478	0	2,764	103	224	32	0	14	0	3,137	4,615	40
2009 Totals	121,592	1,783	123,375	10,923	5,035	15,958	70	15,048	3	196,544	318,136	907

2009 New Jersey Mosquito Trap Count Totals by Week – State Parks

Week of	Male	Female									Total mosquitoes	Sites reporting out of 12
		Anopheles	Aedes	Aedes vexans	Culex	Culex tarsalis	Culex salinarius	Culiseta	Other	Total female		
May 25 - June 1	34	0	395	1	6	2	0	9	0	413	447	6
June 1-8	34	3	417	0	7	0	0	2	0	429	463	7
June 9-14	209	15	383	31	267	0	0	77	0	773	982	9
June 15-21	342	0	565	89	195	12	0	80	0	941	1,283	10
June 22-28	179	1	134	3	31	6	1	102	0	278	457	8
June 29 - July 5	2,392	13	1,954	0	9	84	0	199	0	2,259	4,651	10
July 6-12	656	5	720	32	0	244	0	310	0	1,311	1,967	9
July 13-19	365	0	694	4	0	391	0	409	0	1,498	1,863	7
July 20-26	3,449	24	2,675	932	74	189	0	57	0	3,951	7,400	9
July 27- August 3	421	16	2,599	80	4	134	0	139	0	2,972	3,393	9
August 3-9	203	6	2,263	105	2	32	0	29	0	2,437	2,640	8
August 10-16	193	68	1,520	0	0	25	0	18	0	1,631	1,824	7
August 17-24	44	0	32	0	8	0	0	0	0	40	84	6
August 24-31	48	0	104	16	0	0	0	0	0	120	168	10
August 31- September 7	108	0	1,820	180	16	0	0	0	0	2,016	2,124	7
2009 Totals	8,677	151	16,275	1,473	619	1,119	1	1,431	0	21,069	29,746	122

2009 New Jersey Mosquito Trap Count Totals by Week— Wildlife Refuges													
Week of	Male	Female									Total mosquitoes	Sites reporting out of 3	
		Anopheles	Aedes	Aedes vexans	Culex	Culex tarsalis	Culex salinarius	Culiseta	Other	Total female			
May 25 - June 1	7	0	32	0	2	0	0	0	0	0	34	41	2
June 1-8	32	0	67	0	0	0	0	0	0	0	67	99	2
June 9-14	3	0	1	0	0	0	0	0	0	0	1	4	2
June 15-21	19	0	50	0	16	0	0	0	0	0	66	85	3
June 22-28	43	2	34	0	0	1	0	5	0	0	42	85	3
June 29 - July	33	4	241	0	0	0	0	338	0	0	583	616	3
July 6-12	119	6	206	0	2	24	0	26	0	0	264	383	2
July 13-19	916	100	512	0	0	104	0	692	0	0	1,408	2,324	3
July 20-26	435	165	338	32	16	18	0	106	0	0	675	1,110	3
July 27- August 3	148	78	82	8	24	54	0	93	0	0	339	487	3
August 3-9	160	16	72	8	0	8	0	72	0	0	176	336	2
August 10-16	461	87	386	16	30	61	0	16	0	0	596	1,057	3
August 17-24	0	0	32	0	8	0	0	0	0	0	40	40	2
August 24-31	0	0	4	0	0	0	0	0	0	0	4	4	1
August 31- September 7	0	0	3	0	0	0	0	0	0	0	3	3	1
2009 Totals	2,376	458	2,060	64	98	270	0	1,348	0	0	4,298	6,674	35

Arbovirus Information

More than 2,500 different species of mosquitoes are found worldwide, with about 200 species in the United States. The most common vector in the spread of arboviruses is the mosquito; however, not all mosquitoes are vectors in the transmission of arboviruses.

Male mosquitoes feed almost exclusively on nectar and therefore do not bite. Female mosquitoes lay eggs that require a blood meal and bite animals, warm or cold-blooded, and birds. Stimuli that influence biting include a combination of carbon dioxide, temperature, moisture, smell, color and movement. Humans are seldom the first or second choice for a blood meal. Horses, cattle, smaller mammals and birds are preferred. Although acquiring a blood meal is essential for female egg production, both male and female mosquitoes are mainly nectar feeders.

Mosquito-borne diseases cause more than one million human deaths every year. Some of these diseases include protozoan infections such as malaria; filarial pathogens such as canine heartworm; and viruses that cause dengue, yellow fever and encephalitis.

Arthropod-borne viruses (arboviruses) are the most diverse and serious diseases transmitted to susceptible vertebrate hosts by mosquitoes. All arboviral encephalitides are zoonotic involving a nonhuman primary vertebrate and a primary arthropod vector. Humans and domestic animals can develop clinical illness but usually are “dead-end” hosts because they do not contribute to the transmission cycle.

West Nile virus (WNV) is the most recently emerged arbovirus in North America. West Nile virus is named after the West Nile region of Uganda where it was first discovered in 1937. *Culex* species of mosquitoes are the primary vectors. Common in many parts of the world, WNV had not been seen in the United States until late summer 1999, when it made its debut in New York. WNV then proceeded to travel westward across the continent the following year. West Nile fever can be characterized by fever, headache and rash to more serious symptoms. Although only a small percentage of people infected with WNV display symptoms, WNV can cause encephalitis (an inflammation of the brain) and meningitis (inflammation of the brain and spinal cord) in humans and animals.

Western equine encephalitis (WEE) is mostly found in states west of the Mississippi River. The primary vector is *Culex tarsalis*. Birds are the most important host. Since 1964, there have been fewer than 1,000 cases reported. Human mortality rates are about 5 percent, with horse mortality rates considerably higher.

Eastern equine encephalitis (EEE) is spread to horses and humans by infected mosquitoes. Annually, there are a small number of cases nationwide. EEE is the most serious of the arboviruses that can affect the central nervous system (CNS), resulting in severe complications and even death. Symptoms may range from none at all to flu-like to more serious infections with sudden fever and severe headache followed by seizures and coma. About half of patients die, and of those who survive, many suffer permanent CNS damage.

St. Louis encephalitis (SLE) is transmitted from birds to mammals by an infected mosquito. SLE was discovered in 1933 in St. Louis, Mo. Since then, SLE has been reported in 46 states. Most infections of SLE do not result in illness, with mild cases exhibiting aseptic meningitis or fever. The elderly and very young children are more susceptible, with fatality rates from 2 percent to 20 percent and neurologic dysfunction occurring in about 1 percent of survivors.

The California serogroup is a group of several related viruses that included California encephalitis, La Crosse encephalitis, and Jamestown Canyon virus. Each year, about 75 cases are reported in the United States, with the majority of the illnesses resulting from La Crosse encephalitis. The California serogroup viruses primarily affect male children younger than 16. Infections are mild, with a mortality rate of about four deaths per 1,000 infections.

**North Dakota Mosquito Surveillance
Risk Assessment Chart
for Arbovirus Activity**

Risk Category	Probability of Human Outbreak	Definition of Conditions	Recommended Response by Mosquito Surveillance Team and North Dakota Vector Control Personnel
1a	Remote	Mid-season; first week of July; no observed epizootic activity; low population counts of vector species from New Jersey trap network	Begin preliminary, low-intensity CDC live-trapping network and testing in all areas of the state; test for targeted virus presence.
1b		Late-season; third week of July through September; no observed epizootic activity; high population counts from New Jersey trap network	Deploy mid-intensity CDC live-trapping network and viral testing in areas with high population counts of targeted vector species; continue low intensity trapping and testing in other areas.
2	Low	Sporadic epizootic activity in birds or mosquitoes	Deploy high intensity CDC live-trapping network and viral testing in epizootic areas, and consider preliminary control measures such as source reduction and larval control; continue surveillance in other areas.
3	Moderate	Initial confirmation of virus in horse or human; moderate activity in birds or mosquitoes	Continue as in Category 2; consider adult mosquito-control as indicated by surveillance activity.
4	High	Measures suggesting high risk of human infection (for example, high dead bird densities, high mosquito infection rates, multiple positive mosquito species, horse or mammal cases indicating escalating epizootic transmission, or a human case)	Response as in Category 3; initiate adult mosquito control program in areas of potential human risk.
5	Outbreak in progress	Multiple confirmed human cases; conditions as listed in Category 4	Implement emergency adult mosquito-control program; if widespread, consider aerial spraying.

Appendix A

New Jersey Mosquito Trap Data Analysis

The mosquito's life cycle has four separate and distinct stages: egg, larva, pupa and adult. A female mosquito breeds in the presence of water and lays fertile eggs after obtaining a blood meal. The location in which a female mosquito deposits her eggs in the environment depends upon larval habitat preference. The 36 mosquito species indigenous to North Dakota can be grouped into four categories that reflect their larval habitat preference. These categories include the permanent pool group, the transient water group, the floodwater group, and the artificial container and tree-hole group.

Mosquitoes within the **permanent pool group**, *Anopheles* and *Culex* species, lay eggs either singly or side by side on the water surface of permanent ponds and lakes. Permanent pool mosquitoes can develop continuously in warm water and hatch daily into adults. **Transient water mosquitoes** such as *Culex tarsalis* prefer to lay their eggs in pools of a temporary nature. Common habitats of the transient water group are roadside ditches, canals, ground pools and irrigated lands. Transient water mosquito eggs in ditches and small depressions must wait until rainfall to begin the hatching process. **Floodwater mosquitoes**, the *Aedes* species, lay eggs singly on damp soil or along vegetated shorelines; the eggs remain dormant until these areas are flooded. Once flooded, the eggs hatch if conditions are favorable. Large numbers of larvae emerge, and adults can appear as early as six days after flooding. A major rainstorm, a series of showers, or irrigation sufficient enough to produce standing water promotes hatching in the floodwater species of mosquitoes. The **artificial container and tree-hole group of mosquitoes** place their eggs inside the wall of a container or depression inside a tree, at or above the water line, and the eggs hatch when the water levels rise. A heavy rain resulting in standing water in old tires, tin cans and flowerpots will begin the hatching process for artificial container mosquitoes.

Once hatched, larvae of all species emerge and live in water. After four stages, or instars, the larva molts into a pupa. The pupa stage is a resting, non-feeding stage where the pupa is encased until the adult matures and emerges from the skin after one-and-a-half to four days. Adult male mosquitoes hatch first and live from six to seven days. Female mosquitoes can live for about two weeks but have been found to survive for up to five months with ample food. Peak adult mosquito populations usually appear within two weeks after a number of eggs hatch.

Along with increased rainfall, warmer water temperatures speed up hatching and larval development. If outdoor temperatures are 50 degrees Fahrenheit or higher, productive breeding sites readily produce mosquito larvae. With increasing water temperatures, large mosquito populations can emerge within one week. Research in laboratory settings has shown that if the water temperature exceeds 100 degrees Fahrenheit, it takes only three to four days for larval metamorphosis; if the temperature is 90 degrees Fahrenheit, it takes five days; and a lower water temperature of 70 degrees Fahrenheit decreases rate of growth to 10 days. Floodwater species of *Aedes* larvae generally metamorphose within five to seven days after hatching. The species *Culex tarsalis* completes its life cycle in 14 days at 70 degrees Fahrenheit and in only 10 days at 80 degrees Fahrenheit. On the other hand, some species have naturally adapted to go through their entire life cycle in as little as four days or as long as one month.



When a mosquito becomes an adult, the weather elements affect its peak activity. Most mosquitoes are active from dusk until dawn when wind speeds are less than eight miles per hour, the air temperature is between 65 degrees Fahrenheit and 80 degrees Fahrenheit, and the weather is moderate.

Heavy rains, gusting winds, and cool or high daytime temperatures all limit a mosquito's feeding activity. At temperatures lower than 50 degrees Fahrenheit, mosquitoes become sluggish, reducing their host-seeking behavior. At higher temperatures, usually during daytime hours, adult mosquitoes seek cover in vegetated or humid areas with shade.

Region I

North Dakota Mosquito Surveillance



New Jersey Mosquito Trap Data Analysis





Region II

North Dakota Mosquito Surveillance



New Jersey Mosquito Trap Data Analysis





Region III North Dakota Mosquito Surveillance New Jersey Mosquito Trap Data Analysis



Region IV North Dakota Mosquito Surveillance New Jersey Mosquito Trap Analysis





Region V North Dakota Mosquito Surveillance New Jersey Mosquito Trap Analysis



Region VI

North Dakota Mosquito Surveillance



New Jersey Mosquito Trap Data Analysis



Region VII

North Dakota Mosquito Surveillance



New Jersey Mosquito Trap Data Analysis



Region VIII

North Dakota Mosquito Surveillance



New Jersey Mosquito Trap Data Analysis





State Parks

North Dakota Mosquito Surveillance

New Jersey Mosquito Trap Data Analysis




National Wildlife Refuges North Dakota Mosquito Surveillance New Jersey Mosquito Trap Data Analysis







Appendix B 1994-2009



Weekly New Jersey Mosquito Trap Counts Comparison



Appendix B includes graphs of the annual trap counts from the last week of May through the first week of September. These graphs depict how the mosquito trap counts have changed between 1994 and 2009. Each year, the general trend of North Dakota's mosquito population is a steady rise in population peaking in early to late July, followed by a gradual decrease through the rest of the mosquito season. Yearly and weekly variances in trap numbers can be attributed to factors such as rainfall, temperature and wind speed.




















Weekly Mean of Trap Counts
Sixth Week of August - First Week of September



References

1. American Mosquito Control Association. (1990). *Organization for Mosquito Control* (16-17). Baltimore, Maryland.
2. Darsie, R.F., Jr. and Ward, R.A. (1981). *The Identification and Geographical Distribution of the Mosquitoes of North America North of Mexico*. Fresno, California: American Mosquito Control Association.
3. Moore, C.G. and Pratt, H.D. (1993). *Mosquitoes of Public Health Importance and Their Control* (1-43). Atlanta, GA: Centers for Disease Control and Prevention.
4. Cross, Michael N. (1994, 1995, 1996, 1997, and 1998, September) *1994, 1995, 1996, 1997, and 1998 North Dakota Arbovirus/Mosquito Surveillance Program Annual Reports*. Bismarck, North Dakota: North Dakota Department of Health, Division of Microbiology.
5. Nies, Nicole. (2000). *2000 North Dakota Mosquito Surveillance Program Report*. Bismarck, North Dakota: North Dakota Department of Health, Division of Microbiology.
6. Irmen, Chris and Tescher, Todd. (2001, August). *2001 North Dakota Mosquito Surveillance Program Report*. Bismarck, North Dakota: North Dakota Department of Health, Division of Microbiology.
7. Bold, Tyler and Schulz, Travis. (2002, October). *2002 North Dakota Mosquito Surveillance Program Report*. Bismarck, North Dakota: North Dakota Public Health Laboratory.
8. Schulz, Travis; Bichler, Mike; Voller, Laura; Hoff, Jennifer. (2003, December). *2003 North Dakota Mosquito Surveillance Program Report*. Bismarck, North Dakota: North Dakota Department of Health, Division of Microbiology.
9. Goddard, Laura, et. al. (December 2002). *Vector Competence of California Mosquitoes for West Nile virus*. *Emerging Infectious Diseases* (Vol. 8, No. 12) (1385-1391). University of California, Davis, California.
10. Henke, Michelle; Iverson, Sara; Auen, Cari. (2008). *2008 North Dakota Mosquito Surveillance Program Report*. Bismarck, North Dakota: North Dakota Department of Health, Division of Laboratory Services — Microbiology.