

Vaccines For VFC* and Non-VFC (Insured) Children
(regardless of state of residence)

Vaccine	Ages Covered	Eligibility Criteria
Hepatitis B	Birth	<ul style="list-style-type: none"> • <u>Available for administration by hospitals to all children (VFC and insured).</u> • The ACIP and the NDDoH strongly recommend that all children receive the birth dose of hepatitis B vaccine. • Hepatitis B vaccine is VFC-only for all other doses in the hepatitis B series.

Vaccines For VFC Children Only
(regardless of state of residence)

Vaccine	Ages Covered	Eligibility Criteria
DT	6 weeks – 6 years	<ul style="list-style-type: none"> • Only to be administered when DTaP is medically contraindicated. • Call for approval prior to ordering.
DTaP	6 weeks – 6 years	
DTaP/HepB/IPV (Pediarix [®])	6 weeks – 6 years	<ul style="list-style-type: none"> • Approved only for the primary series (i.e., 2, 4, and 6 months). • <u>Only available for administration by IHS facilities or other facilities serving significant American Indian populations.</u>
DTaP-IPV/Hib (Pentacel [®])	6 weeks – 59 months	<ul style="list-style-type: none"> • Approved for use at 2, 4, 6, and 12 – 18 months of age • Pediarix[®] and PedvaxHIB[®] preferred for IHS and other facilities serving significant American Indian populations.
DTaP-IPV (Kinrix [®])	4 years – 6 years	Approved for use as the fifth dose of DTaP and fourth dose of IPV at 4 – 6 years of age.
Hepatitis A	1 – 18 years	Should be routinely administered to children 12-23 months of age.
Hepatitis B	Birth – 18 years	
Hib	6 weeks – 59 months	PedvaxHIB [®] is a three-dose Hib vaccine administered at 2, 4, and 12 – 15 months of age.
HPV4 (Gardasil [®])	9 years – 18 years	Available for both males and females.
Influenza	6 months – 18 years	
IPV	6 weeks – 18 years	
Meningococcal Conjugate (MCV4, Menactra [®])	2 – 18 years	Should be routinely administered to children 11-12 years of age, with a booster dose at 16 years of age.
MMR	12 months – 18 years	
Pneumococcal Conjugate (PCV13, Prevnar13 [®])	6 weeks – 59 months	The ACIP recommends PCV13 for all children ages two through 59 months and for children ages 60 through 71 months who have underlying medical conditions that increase their risk of pneumococcal disease or

		complications. Children who were completely vaccinated with PCV7, need a supplemental dose of PCV13.
Pneumococcal Polysaccharide	2 – 18 years	<ul style="list-style-type: none"> • Available for high-risk children with one or more of the following conditions: <ul style="list-style-type: none"> ○ Chronic illnesses such as cardiovascular disease, chronic pulmonary disease, diabetes mellitus, alcoholism, chronic liver disease, or CSF leaks. ○ Functional or anatomic asplenia (splenectomy) ○ Conditions associated with immunosuppression, including HIV infection, leukemia, lymphoma, Hodgkin’s disease, multiple myeloma, generalized malignancy, chronic renal failure, nephritic syndrome, or other conditions associated with immunosuppression (organ or bone marrow transplantation); and people receiving immunosuppressive chemotherapy, including long-term high-dose corticosteroids. ○ Cochlear implants
Rotavirus	6 weeks – 8 months, 0 days	Both Rotateq [®] (three dose rotavirus vaccine to be administered at 2, 4, and 6 months of age) and Rotarix [®] (two dose rotavirus vaccine to be administered at 2 and 4 months of age) are available.
Td	7 – 18 years	<ul style="list-style-type: none"> • Tdap is recommended instead of Td for adolescents and adults. • Td is only necessary for children and adolescents who have not completed the primary series of DTaP.
Tdap	7 – 18 years	<ul style="list-style-type: none"> • Tdap is recommended instead of Td for adolescents and adults. • Boostrix[®] and Adacel[®] are available for order.
Varicella (Chickenpox)	12 months – 18 years	
Other State-Supplied Vaccines (regardless of state of residence)		
Vaccine	Ages Covered	Eligibility Criteria
HPV4 (Gardasil [®])	19 – 26 years	<p>Available for uninsured and underinsured males and females</p> <ul style="list-style-type: none"> • Adults ages 19 – 21 years who are enrolled in Medicaid must be vaccinated with private vaccine and Medicaid should be billed. • Adults ages 22 – 26 years who are enrolled in Medicaid may be given state-supplied vaccine. • Insured individuals must be vaccinated with private vaccine and insurance should be billed.
Meningococcal Conjugate (MCV-4, Menactra [®])+	17 years – 55 years	<ul style="list-style-type: none"> • Available for all college freshmen attending colleges/universities, including those with health insurance. • Also available for all high school seniors, including those with health insurance.

MMR	19 years and older	Available to <u>uninsured</u> or <u>underinsured</u> students born in or after 1957 enrolled in a college/university.
Pneumococcal Polysaccharide	19 years and older	Available for high-risk <u>uninsured</u> or <u>underinsured</u> adults, with one or more of the following conditions: <ul style="list-style-type: none"> • Chronic illnesses such as cardiovascular disease, chronic pulmonary disease (including asthma), diabetes mellitus, alcoholism, chronic liver disease, or CSF leaks. • Functional or anatomic asplenia (splenectomy) • Conditions associated with immunosuppression, including HIV infection, leukemia, lymphoma, Hodgkin’s disease, multiple myeloma, generalized malignancy, chronic renal failure, nephritic syndrome, or other conditions associated with immunosuppression (organ or bone marrow transplantation); and people receiving immunosuppressive chemotherapy, including long-term high-dose corticosteroids. • Smokers
Td/Tdap+	≥ 7 years	<ul style="list-style-type: none"> • Available for all (including those with health insurance) individuals who have or who anticipate having close contact with an infant aged <12 months: <ul style="list-style-type: none"> ○ Parents/guardians of infants < 12 months ○ Childcare providers, regardless of age of children attending childcare ○ Expecting fathers • Available for any uninsured or underinsured adult. • Tdap is recommended instead of Td for adolescents and adults.

+The NDDoH also supplies MCV4 and Tdap to local public health units to vaccinate insured children for the middle school entry immunization requirements.

*VFC-eligible children are those who are 18 and younger and meet one of the following criteria:

- No health insurance
- Medicaid eligible
- American Indian or Alaskan Native
- Underinsured – have health insurance, but it does not cover a particular vaccine
 - If your clinic does not have an agreement with a federally qualified health center to vaccinate underinsured children on their behalf, then your clinic must refer underinsured children to a rural health center or federally qualified health center to receive VFC vaccine.

Updates since the previous version of the coverage table are shown in red.