USGS
science for a changing world

An Improved Understanding of Nitrogen and Phosphorus Delivery : Results from Refined Regional SPARROW Models

By
Joel M. Galloway
Dale M. Robertson[1], David A. Saad[1], Gregory E. Schwarz[2], and Richard A. Alexander[2]
U.S. Geological Survey,
[1]Wisconsin Water Science Center; [2]National Center, Reston

U.S. Department of the Interior
U.S. Geological Survey

Approach - SPARROW Water-Quality Model -

SPAtially Referenced Regression on Watershed Attributes

http://water.usgs.gov/nawqa/sparrow



- Mass Balance Model with spatially variable deliveries. Hybrid statistical/ mechanistic process structure. Data-driven, nonlinear estimation of parameters
- Separates land and in-stream processes
- Predictions of mean-annual flux reflect long-term, net effects of nutrient supply and loss processes in watersheds
- Once calibrated, the model has physically interpretable coefficients; model supports hypothesis testing and uncertainty estimation

Watershed Modeling Continuum





Cole et al. 1997



SWAT Users Manual

MO/MSSP SPARROW Model Calibration

One Source: 2002 Point (WWTP) TN inputs, kg



9,182 WWTPs

One Land-to-Water Delivery: Tile Drains



Calibration



24,475 Catchments based on RF1 River Network



Long-term detrended Loads for 856 sites

Delivered Incremental Yields

Nitrogen



Phosphorus



Robertson and Saad, 2013

Ranking of State Contributions to the Gulf of Mexico from the MARB



Robertson et al., 2014

Ranking of State Contributions to the Gulf of Mexico from the MARB



Robertson et al., 2014

Regional Models part of the NAWQA Program

MRB SPARROW
Lead Scientists
Coordinator – Steve Preston

Robertson & Saad, WI

2002 SPARROW Models

All Published in 2011 in JAWRA

Distribution in Incremental Phosphorus Yields



Robertson and Saad, 2011

Relative Importance of Various Phosphorus Sources – Spatially Explicit











Robertson and Saad, 2011

Binational SPARROW Model for the Upper Midwest



G. Benoy (IJC), W. Jenkinson (NRC-Canada), C. Johnston (USGS)

USGS science for a changing world

NRC·CNRC

Methods to demonstrate results and help guide decisions > Nutrient Reduction Strategies

1. ***SPARROW Mapper –***

   Easy and simple way to get SPARROW results, especially by hydrologic and political boundaries.

http://wim.usgs.gov/SparrowMRB3/SparrowMRB3Mapper.html#

http://wim.usgs.gov/SparrowGL/SparrowGLMapper.html#

http://wim.usgs.gov/SparrowMARB/SparrowMARBMapper.html#

MARB SPARROW MAPPER




USGS
science for a changing world
SPARROW Mapper
Mississippi Basin 2002 Nutrient Models
Links
Zoom History
Available Layers
Nutrient model results
Total Phosphorus
Total Nitrogen
Area of Interest
North Dakota
Select Main Watershed
Select HUC8
Select Tributary Outlet
Displayed Metric
Delivered Accumulated Yield
2002 Total Phosphorus Delivered Accumulated Yield (kg/km²)
Delivered Accumulated Yield (kg/km²)
Ranked by HUC
Forest/Wetland
Loess
Stream Channel
Manure (Confined and Unconfined)
Farm Fertilizer
Urban Land
Sewerage Point Sources
Export Data
Delivered Accumulated Yield for entire view
View Nutrient Totals
Lake Manitoba
Winnipeg
Selkirk
Portage la Prairie
Brandon
Weyburn
Estevan
Steinbach
Minot
Wolf Point
Grand Forks
GREAT PLAINS
SOUTH DAKOTA
Pierre
Rapid City
Gillette
Badlands
200 km
100 mi


USGS
science for a changing world
SPARROW Mapper
Mississippi Basin 2002 Nutrient Models
Links
Zoom History
Available Layers
Nutrient model results
Total Phosphorus
Total Nitrogen
Area of Interest
North Dakota
Select Main Watershed
Select HUC8
Select Tributary Outlet
Displayed Metric
2002 Total Phosphorus Delivered Accumulated Yield (kg/km²)
Delivered Accumulated Yield (kg/km²)
14
12
10
8
6
4
2
0
1
2
Ranked by HUC
Forest/Wetland
Loess
Stream Channel
Manure (Confined and Unconfined
Farm Fertilizer
Urban Land
Sewerage Point Sources
Export Data
Delivered Accumulated Yield for entire view
View Nutrient Totals
Lake Manitoba
Winnipeg
Selkirk
Portage la Prairie
Brandon
Weyburn
Souris
Red
Steinbach
Estevan
Minot
Wolf Point
Grand Forks
GREAT PLAINS
SOUTH DAKOTA
Pierre
Gillette
Rapid City
Badlands
200 km
100 mi

MRB3 SPARROW Mapper for Delivered P Yields – shown by Major Watershed




http://wim.usgs.gov/SparrowMRB3/SparrowMRB3mapper.html
Sparrow MRB3 Mapper
File Edit View Favorites Tools Help
USGS science for a changing world
SPARROW Mapper
MRB3 2002 Nutrient Models
Links
Zoom History
ONTARIO
Winnipeg
Brandon
Kenora
Nipigon
Thunder Bay
Wawa
Timmins
Matagami
Rouyn-Noranda
Val-d'Or
Sudbury
Sault Ste. Marie
Lake Superior
BADLANDS
NORTH DAKOTA
Bismarck
GREAT PLAINS
SOUTH DAKOTA
Pierre
Lake Michigan
Lake Huron
Georgian Bay
Owen Sound
Barrie
Kingston
Toronto
Lake Ontario
Hamilton
London
Sarnia
Lake Erie
Milwaukee
Chicago
Washington
Richmond
VIRGINIA
Norfolk
Knoxville
Greensboro
Raleigh
NORTH CAROLINA
Memphis
Available Layers
Nutrient model results
Total Phosphorus
Total Nitrogen
Area of Interest
Select State
Select Major Watershed
Select HUC
Select Tributary
Mapped Metric
Delivered Accumulate...
Group results by
Major Watersheds
2002 Total Phosphorus Delivered Accumulated Load (kg) by Major Watersheds
0. - 122890
122890 - 691535
691535 - 2109061
2109061 - 4714617
4714617 - 31292801
Additional Layers
MRB3 Area
Stream reaches (RF1)
Main Stems
SPARROW calibration sites
2002 Total Phosphorus Delivered Accumulated Load (kg) by Major Watersheds
Upper Miss #1
Red R #7
Souris R #13
Delivered Accumulated Load (kg)
Ranked by Major Watershed
Export Data
Delivered Accumulated Load for entire view
View Nutrient Sources
100%


SPARROW Mapper
MRB3 2002 Nutrient Models
Links
Zoom History
Available Layers
Nutrient model results
Total Phosphorus
Total Nitrogen
Area of Interest
Select State
Select Major Watershed
Select HUC
Select Tributary
Mapped Metric
Delivered Accumulate...
Group results by
Major Watersheds
2002 Total Phosphorus Delivered Accumulated Load (kg) by Major Watersheds
0. - 122890
122890 - 691535
691535 - 2109061
2109061 - 4714617
4714617 - 31292801
Additional Layers
MRB3 Area
Stream reaches (RF1)
Main Stems
SPARROW calibration sites
2002 Total Phosphorus Delivered Accumulated Load (kg) by Major Watersheds
Forest/Wetland
Manure (Unconfined)
Manure (Confined)
Farm Fertilizer
Urban Land
Point Sources
Delivered Accumulated Load (kg)
Red River
Major Watershed
Export Data
Delivered Accumulated Load for entire view
View Nutrient Totals
ONTARIO
NORTH DAKOTA
SOUTH DAKOTA
GREAT PLAINS
Winnipeg
Brandon
Bismarck
Pierre
Thunder Bay
Sault Ste. Marie
Sudbury
Timmins
Toronto
Hamilton
London
Milwaukee
Chicago
Detroit
Windsor
Sarnia
Owen Sound
Barrie
Kingston
Washington
Richmond
Norfolk
Raleigh
Greensboro
Knoxville
Memphis
Lake Superior
Lake Michigan
Lake Huron
Lake Erie
Lake Ontario
Georgian Bay
Algonquin Provincial Park


USGS
science for a changing world
SPARROW Mapper
MRB3 2002 Nutrient Models
Links
Zoom History
Available Layers
Nutrient model results
Total Phosphorus
Total Nitrogen
Area of Interest
Select State
Select Major Watershed
Select HUC
Select Tributary
Mapped Metric
Delivered Accumulate...
Group results by
Major Watersheds
2002 Total Phosphorus Delivered Accumulated Load (kg) by Major Watersheds
0. - 122890
122890 - 691535
691535 - 2109061
2109061 - 4714617
4714617 - 31292801
Additional Layers
MRB3 Area
Stream reaches (RF1)
Main Stems
SPARROW calibration sites
2002 Total Phosphorus Delivered Accumulated Load (kg) by Major Watersheds
Forest/Wetland
Manure (Unconfined)
Manure (Confined)
Farm Fertilizer
Urban Land
Point Sources
Delivered Accumulated Load (kg)
140000
120000
100000
80000
60000
40000
20000
0
Souris River
Major Watershed
Export Data
Delivered Accumulated Load for entire view
View Nutrient Totals
ONTARIO
Winnipeg
Brandon
Kenora
Nipigon
Thunder Bay
Wawa
Timmins
Matagami
Rouyn-Noranda
Val-d'Or
Sudbury
Sault Ste. Marie
Lake Superior
BADLANDS
NORTH DAKOTA
Bismarck
GREAT PLAINS
SOUTH DAKOTA
Pierre
Lake Michigan
Lake Huron
Georgian Bay
Owen Sound
Barrie
Kingston
Toronto
Hamilton
London
Sarnia
Lake Erie
Milwaukee
Chicago
Washington
Richmond
Norfolk
VIRGINIA
Knoxville
Greensboro
Raleigh
NORTH CAROLINA
TENNESSEE
Memphis
Algonquin Provincial Park
100%

Methods to demonstrate results and help guide decisions

2. ***Decision Support System*** ***Scientists/Managers – Capable of using to visualize SPARROW output and run various scenarios.***

http://cida.usgs.gov/sparrow/

Booth et al., 2011

SPARROW Decision Support System

Find a Model by Geographic Location:

Select a region or state. When a state is selected, all models containing that state are listed.



Iowa

Find a Model by Modeled Constituent:

Any

Models matching your criteria (click a model to show details)

- Mississippi/Atchafalaya Basin Total Nitrogen Model - 2002
- Mississippi/Atchafalaya Basin Total Phosphorus Model - 2002
- National Suspended Sediment Model - 1992
- National Total Nitrogen Model - 1992
- National Total Organic Carbon Model
- National Total Phosphorus Model - 1992
- Total Nitrogen Model for the Great Lakes, Ohio, Upper Mississippi, and Souris-Red-Rainy Region - 2002
- Total Nitrogen Model for the Missouri River Basin - 2002
- Total Phosphorus Model for the Great Lakes, Ohio, Upper Mississippi, and Souris-Red-Rainy Region - 2002
- Total Phosphorus Model for the Missouri River Basin - 2002

Documentation and Further Reading

- What is SPARROW?
- What is SPARROW Decision Support?
- SPARROW Applications & Documentation
- SPARROW DSS FAQs

Tutorial Videos

Select a video...

Watch now >>

Found a bug or have a comment?

Please send bugs, suggestions and questions to the SPARROW Decision Support System Administrator.

Selected Model

Mississippi/Atchafalaya Basin Total Phosphorus Model - 2002

Explore this model in the Decision Support System >>

Modeled Constituent:	Phosphorus
Base Year:	2002
Stream Network:	Enhanced River Reach File 2.0 *Geometry and additional reach and network attribute data are available with the stream network data, which is available as a separate download.*
Model Updates:	View this model's updates

Watershed Based Sessions

To start the DSS with the outlet river reach of a major watershed selected for downstream tracking, select a watershed and click *Go*.

Go >>

Scenario Based Sessions

To start the DSS with a predefined scenario, click on the link for one of the scenarios below.


USGS
science for a changing world
USGS Home
Contact USGS
Search USGS
SPARROW Decision Support System Mississippi/Atchafalaya Basin Total Phosphorus Model - 2002
<< Home
Display Results
Downstream Tracking
Change Inputs
Map the effect of management scenarios on stream water quality, based on hypothetical changes in source inputs. For more information, click here.
1. Select stream reach(es) where changes will be applied
--Locate on map
--Find by name or hydrologic unit code
2. Change the values of the source inputs
(Right click to change input values or show on map)
Missouri River
upstream of 29040 (REACH:29040)
3. Display Results
From the Display Results tab, select a data series.
(Map relative or absolute changes using the Comparison to Original Model feature)
Map settings have changed. Update map to refresh all data.
Update Map
Find a reach...
Export Data...
Session
Layers
Hide Header/Footer
SPARROW Model / Videos
Remove overlay
Missouri River
Input Changes
Notes
Wastewater Treatment Plants
Phosphorus (kg•year⁻¹)
1
enter custom multiplier...
Manure (total)
Phosphorus (kg•year⁻¹)
0.5
enter custom multiplier...
Fertilizers (farm)
Phosphorus (kg•year⁻¹)
0.5
enter custom multiplier...
Forest and Wetland Areas
Land Area (km²)
1
enter custom multiplier...
Urban Areas
Land Area (km²)
1
enter custom multiplier...
Channel Erosion
Channel Length (m)
1
enter custom multiplier...
Save Group Changes
Cancel
EXPLANATION
Incremental Load
kg•year⁻¹ of Phosphorus
< 300
300 to 1100
1100 to 3000
3000 to 8500
> 8500
USGS
50 mi.
100 km.
Map out of sync


Display Results
Downstream Tracking
Change Inputs
Map the model results by reach or catchment.
1. Select a Data Series
Data Series
Incremental Yield
Comparison To Original Model
% change from original
2. Select a Model Source
Model Source
All
Map Units: Mass Percent
3. Select the map display options
Display: Reaches Catchments
Calibration Sites
Reach Overlay
HUC8 Overlay
Binning for Map Color and Legend
5 Equal Count Bins
Auto binning
Edit Custom Bins...
Currently mapping Change in Incremental Yield.
The map is up to date.
Find a reach...
Export Data...
Session
Layers
Remove overlay
Show Header/Footer
SPARROW Model / Videos
EXPLANATION
Change in Incremental Yield
Percentage of Phosphorus
-50 to -38
-38 to -28
-28 to -11
-11 to -1
-1 to 0
USGS

MISSOURI R (ID: 29040)

Reach/Catchment Info | Model Source Inputs | Predicted Values | **Graphs**

Sources

Orig.	Adj.		Orig.	Adj.	
		Wastewater Treatment Plants			Manure (total)
		Fertilizers (farm)			Forest and Wetland Areas
		Urban Areas			Channel Erosion
		Loess Soils			

Total Phosphorus Load by Source (kg year⁻¹)



Share of Total Phosphorus Load by Source - Original



Share of Total Phosphorus Load by Source - Adjusted



OK | Cancel

MISSOURI R (ID: 29040)

Reach/Catchment Info | Model Source Inputs | **Predicted Values** | Graphs

Current Mapped Value: 7.14 kg•km^{-2}•yr^{-1} of Phosphorus (Incremental Yield)

Predicted Values (Data Series)

Source	Original (Phosphorus kg•year^{-1})	% of Load (Orig.)	Adjusted (Phosphorus kg•year^{-1})	% of Load(Adj.)	% Change
Total Load					
Wastewater Treatment Plants Total L...	115,077	4.8	115,077	6.2	0
Manure (total) Total Load	573,253	23.8	286,627	15.3	-50
Fertilizers (farm) Total Load	506,760	21.0	253,380	13.6	-50
Forest and Wetland Areas Total Load	235,110	9.8	235,110	12.6	0
Urban Areas Total Load	290,095	12.0	290,095	15.5	0
Channel Erosion Total Load	687,324	28.5	687,324	36.8	0
Loess Soils Total Load	0	0.0	0	0.0	0
Total Load	2,407,620	100.0	1,867,614	100.0	-22
Incremental Load					
Wastewater Treatment Plants Increm...	0	0.0	0	0.0	0
Manure (total) Incremental Load	1,056	48.7	528	39.8	-50
Fertilizers (farm) Incremental Load	629	29.0	314	23.7	-50
Forest and Wetland Areas Increment...	50	2.3	50	3.7	0
Urban Areas Incremental Load	434	20.0	434	32.7	0
Channel Erosion Incremental Load	0	0.0	0	0.0	0
Loess Soils Incremental Load	0	0.0	0	0.0	0
Incremental Load	2,167	100.0	1,325	100.0	-39

OK | Cancel

References:

Robertson, D.M., Saad, D.A., Schwarz, G.E., 2014, Spatial Variability in Nutrient Transport by HUC8, State, and Subbasin Based on Mississippi/Atchafalaya River Basin SPARROW Models: Journal of the American Water Resources Association.

Robertson, D.M. and Saad, D.A., 2013, SPARROW models used to understand nutrient sources in the Mississippi/Atchafalaya River Basin: Journal of Environmental Quality. v. 42, no. 5, p. 1422-1440, DOI: 10.2134/jeq2013.02.0066.

Robertson, D.M. and D.A. Saad, 2011. Nutrient Inputs to the Laurentian Great Lakes by Source and Watershed Estimated Using SPARROW Watershed Models. Journal of the American Water Resources Association. v. 47, p. 1011-1033, DOI: 10.1111/j.1752-1688.2011.00574.x.

Booth, N.L., E.J. Everman, I.-L. Kuo, L. Sprague, and L. Murphy, 2011. A Web-Based Decision Support System for Assessing Regional Water-Quality Conditions and Management Actions. Journal of the American Water Resources Association, v. 47, p. 1136-1150.

Saad, D.A., G.E. Schwarz, D.M. Robertson, and N.L. Booth, 2011. A Multi-Agency Nutrient Dataset Used to Estimate Loads, Improve Monitoring Design, and Calibrate Regional Nutrient SPARROW Models. Journal of the American Water Resources Association, v. 47, p. 933-949, DOI: 10.1111/j.1752-1688.2011.00575.x

Robertson, D.M., Schwarz, G.E., Saad, D.A., and Alexander, R.B., 2009, Incorporating uncertainty into the ranking of SPARROW model nutrient yields from Mississippi/Atchafalaya River basin watersheds. Journal of the American Water Resources Association, v. 45, n. 2, p. 534-549.

Alexander, R.B., Smith, R.A., Schwarz, G.E., Boyer, E.W., Nolan, J.V., and Brakebill, J.W., 2008, Differences in phosphorus and nitrogen delivery to the Gulf of Mexico from the Mississippi River Basin. Environmental Science and Techology, v. 42, n. 3, p. 822-830.