



### **About your medicine:**

Azithromycin is an antibiotic used to treat infections

### **Before using this medicine:**

Tell the healthcare professional if you...

- Have had or are currently ill from kidney or liver disease
- Have had or are currently ill from colitis or stomach problems
- Have ever had a severe allergic reaction to antibiotics; especially erythromycin
- Are breastfeeding

### **Possible side effects of this medicine:**

If these symptoms continue or become severe, notify a healthcare professional...

- Stomach upset
- Loose stools/diarrhea
- Loss of appetite
- Nausea and vomiting
- Stomach cramps
- Vaginal irritation
- Yellow skin or eyes

### **How to take this medicine:**

- This is the only dose of this medicine you will receive
- Take all 4 tablets at once with one glass (8 ounces) of water
- DO NOT take antacids (Maalox, Mylanta, Rolaids, Tums, etc.) for 2 hours after taking this medicine
- Drink several additional glasses of water today

### **Warning:**

Call a healthcare professional immediately if you develop any of the following:

- Breathing difficulties
- Swelling of the face
- Dark urine
- **(DO NOT** take any “over-the-counter” medicines until you talk with a healthcare professional)

### **Sexual Behavior:**

**Abstain from any type of sexual activity for 7 days after treatment of self and partner**

**For more information, contact the NDDoH STD Program at 701.328.2378 or toll-free 800.472.2180**