

HIV, HCV and STD Screening Resource Guide

	HIV	HCV	STD
SEX:			
• Sexual partners of HIV infected persons	Annually /3-6 months	At Least Once in Lifetime	Annually/ 3-6 Months
• Multiple sex partners	Annually		Annually
• Men who have sex with men (MSM)	Annually	Annually	Annually / 3-6 Months
• Having unprotected sex	Annually		Annually
• Unprotected anal sex	Annually		Rectal Testing
• Unprotected oral sex	At Least Once in Lifetime		Pharyngeal Testing
• Persons who exchange sex for money or drugs	Annually /3-6 months	Annually	Annually
• All pregnant women	Test With Each Pregnancy	At-Risk Individuals	Test With Each Pregnancy
Drugs:			
• Used drugs not prescribed to them (non-injection)		At Least Once in Lifetime	
• Current or past Injection drug users	Annually / 3-6 Months	Annually / 3-6 Months	
• Share or have shared needles or have had needle exposure	Annually / 3-6 Months	Annually / 3-6 Months	
Tattoos:			
• Tattoos or body piercings in an unsterile environment	Annually / 3-6 Months	Annually / 3-6 Months	
Diagnosed with Disease:			
• Have HIV or AIDS		At Diagnosis/As Needed	At Diagnosis/Annually
• Persons diagnosed with Tuberculosis	At Diagnosis		
• Persons diagnosed with HCV	At Diagnosis		
• Persons diagnosed with a STD	At Diagnosis		
Received Treatment:			
• Received donated blood or organs before 1992		At Least Once in Lifetime	
• Received blood clotting factors before 1987		At Least Once in Lifetime	
• Are getting long term hemodialysis		At Least Once in Lifetime	
• Undiagnosed liver problems		At Least Once in Lifetime	