

**Vaccines For VFC\* and Non-VFC (Insured) Children**  
**(regardless of state of residence)**

Vaccine	Ages Covered	Eligibility Criteria
Hepatitis B	Birth	<ul style="list-style-type: none"> <li>Available for administration at birth by hospitals to all children (VFC and insured).</li> <li>The ACIP and the NDDoH <i>strongly</i> recommend that all children receive the birth dose of hepatitis B vaccine.</li> </ul>

**Vaccines for VFC\* (regardless of state of residence) and vaccines for insured North Dakota children who are vaccinated at local public health units**

Vaccine	Ages Covered	Eligibility Criteria
DT	6 weeks – 6 years	<ul style="list-style-type: none"> <li>Only to be administered when DTaP is medically contraindicated.</li> <li>Call for approval prior to ordering.</li> </ul>
DTaP	6 weeks – 6 years	Both Daptacel <sup>®</sup> and Infanrix <sup>®</sup> are available.
DTaP/HepB/IPV (Pediarix <sup>®</sup> )	6 weeks – 6 years	Approved only for the primary series (i.e., 2, 4, and 6 months).
DTaP-IPV/Hib (Pentacel <sup>®</sup> )	6 weeks – 4 years	Approved for use at 2, 4, 6, and 12 – 18 months of age
DTaP-IPV (Kinrix <sup>®</sup> )	4 years – 6 years	Approved for use as the fifth dose of DTaP and fourth dose of IPV at 4 – 6 years of age.
Hepatitis A	1 – 18 years	<ul style="list-style-type: none"> <li>Should be routinely administered to children 12-23 months of age.</li> <li>Both Havrix<sup>®</sup> and Vaqta<sup>®</sup> are available.</li> </ul>
Hepatitis B	Birth – 18 years	Both Engerix-B <sup>®</sup> and Recombivax <sup>®</sup> are available.
Hib	6 weeks – 59 months	<ul style="list-style-type: none"> <li>PedvaxHIB<sup>®</sup> is a three-dose series at 2, 4, and 12 – 15 months of age.</li> <li>ActHib<sup>®</sup> is a four-dose series at 2, 4, 6, and 12 – 15 months.</li> <li>Hiberix<sup>®</sup> is approved for booster dose at 12 – 15 months.</li> </ul>
Hepatitis B – Hib (Comvax <sup>®</sup> )	6 weeks – 4 years	<ul style="list-style-type: none"> <li>Approved for use at 2, 4, and 12 – 15 months of age.</li> <li>Cannot be used for the birth dose of hepatitis B vaccine.</li> </ul>
HPV	9 years – 18 years	<ul style="list-style-type: none"> <li>Gardasil<sup>®</sup> is available for both males and females ages 9 and older.</li> <li>Cervarix<sup>®</sup> is available for only females ages 9 and older.</li> </ul>
Influenza	6 months – 18 years	
IPV	6 weeks – 18 years	
Meningococcal Conjugate (MCV4)	2 – 18 years	<ul style="list-style-type: none"> <li>Should be routinely administered to children 11-12 years of age, with a booster dose at 16 years of age.</li> <li>Both Menactra<sup>®</sup> and Menveo<sup>®</sup> are available.</li> </ul>
MMR	12 months – 18 years	
Pneumococcal	6 weeks – 59 months	The ACIP recommends PCV13 for all children ages two

Conjugate (PCV13, Prevnar13 <sup>®</sup> )		through 59 months and for children ages 60 through 71 months who have underlying medical conditions that increase their risk of pneumococcal disease or complications. Children who were completely vaccinated with PCV7 need a supplemental dose of PCV13.
Pneumococcal Polysaccharide (PPV23)	2 – 18 years	<ul style="list-style-type: none"> <li>• Available for high-risk children with one or more of the following conditions: <ul style="list-style-type: none"> <li>○ Chronic illnesses such as cardiovascular disease, chronic pulmonary disease, diabetes mellitus, alcoholism, chronic liver disease or CSF leaks.</li> <li>○ Functional or anatomic asplenia (splenectomy)</li> <li>○ Conditions associated with immunosuppression, including HIV infection, leukemia, lymphoma, Hodgkin’s disease, multiple myeloma, generalized malignancy, chronic renal failure, nephritic syndrome, or other conditions associated with immunosuppression (organ or bone marrow transplantation); and people receiving immunosuppressive chemotherapy, including long-term high-dose corticosteroids.</li> <li>○ Cochlear implants</li> </ul> </li> </ul>
Rotavirus	6 weeks – 8 months, 0 days	Both Rotateq <sup>®</sup> (three dose rotavirus vaccine to be administered at 2, 4, and 6 months of age) and Rotarix <sup>®</sup> (two dose rotavirus vaccine to be administered at 2 and 4 months of age) are available.
Td	7 – 18 years	<ul style="list-style-type: none"> <li>• Tdap is recommended instead of Td for adolescents and adults.</li> <li>• Td is only necessary for children and adolescents who have not completed the primary series of DTaP.</li> <li>• Both Decavac<sup>®</sup> and Massachusetts Biologics are available.</li> </ul>
Tdap	7 – 18 years	<ul style="list-style-type: none"> <li>• Tdap is recommended instead of Td for adolescents and adults.</li> <li>• Boostrix<sup>®</sup> and Adacel<sup>®</sup> are available for order.</li> </ul>
Varicella (Chickenpox)	12 months – 18 years	

**Uninsured/Underinsured Adult Vaccines  
(regardless of state of residence)**

Vaccine	Ages Covered	Eligibility Criteria
HPV	19 – 26 years	<p>Available for uninsured and underinsured males and females.</p> <ul style="list-style-type: none"> <li>• Adults ages 19 – 21 years who are enrolled in Medicaid must be vaccinated with private vaccine and Medicaid should be billed (considered insured).</li> <li>• Adults ages 22 – 26 years who are enrolled in Medicaid may be given state-supplied vaccine (considered underinsured).</li> <li>• Insured individuals must be vaccinated with private vaccine and insurance should be billed.</li> </ul>
Meningococcal	19 years – 55 years	Available for uninsured and underinsured adults.

Conjugate (MCV-4)		
MMR	19 years and older	Available for uninsured and underinsured adults.
Pneumococcal Polysaccharide	19 years and older	Available for high-risk <u>uninsured</u> or <u>underinsured</u> adults, with one or more of the following conditions: <ul style="list-style-type: none"> <li>• Chronic illnesses such as cardiovascular disease, chronic pulmonary disease (<b>including asthma</b>), diabetes mellitus, alcoholism, chronic liver disease, or CSF leaks.</li> <li>• Functional or anatomic asplenia (splenectomy)</li> <li>• Conditions associated with immunosuppression, including HIV infection, leukemia, lymphoma, Hodgkin’s disease, multiple myeloma, generalized malignancy, chronic renal failure, nephritic syndrome, or other conditions associated with immunosuppression (organ or bone marrow transplantation); and people receiving immunosuppressive chemotherapy, including long-term high-dose corticosteroids.</li> <li>• <b>Smokers</b></li> </ul>
Td/Tdap	19 years and older	<ul style="list-style-type: none"> <li>• Available for uninsured and underinsured adults.</li> <li>• A single dose is recommended for all adults, including those 65 years and older.</li> </ul>

\*VFC-eligible children are those who are 18 and younger and meet one of the following criteria:

- No health insurance
- Medicaid eligible
- American Indian or Alaskan Native
- Underinsured – have health insurance, but it does not cover a particular vaccine

NDIIS VFC Eligibility Entry:

- Children 18 and younger:
  - Medicaid
  - American Indian
  - Uninsured
  - Underinsured
  - Other State Eligible – insured children receiving hepatitis B birth dose or insured children vaccinated at universal local public health units
  - Not Eligible – all other insured children
- Adults
  - Other State Eligible – uninsured/underinsured adults receiving state-supplied HPV, Tdap, MMR, MCV4, or PPSV23
  - Not Eligible – all other adults